Royal College of Nursing

Clinical Nurse Specialist in Endometriosis

CLINICAL PROFESSIONAL RESOURCE



Acknowledgements

The RCN would like to thank the project team who originally developed these standards in 2018. This third edition was updated in 2024 by:

Debra Holloway FRCN, retired Nurse Consultant Gynaecology, Guy's and St Thomas' NHS Foundation Trust and RCN Women's Health Forum

Dr Wendy Norton FRCN, Honorary Senior Research Fellow, Faculty of Health and Life Sciences, De Montfort University, Leicester, and RCN Women's Health Forum

This document has been designed in collaboration with our members to ensure it meets most accessibility standards. However, if this does not fit your requirements, please contact corporate.communications@rcn.org.uk

RCN Legal Disclaimer

This publication contains information, advice and guidance to help members of the RCN. It is intended for use within the UK but readers are advised that practices may vary in each country and outside the UK. The information in this booklet has been compiled from professional sources, but its accuracy is not guaranteed. Whilst every effort has been made to ensure the RCN provides

Contents

1 Introduction 4

2 Defining endometriosis 5

3 The role of the clinical nurse specialist (CNS) in endometriosis 6

3.1 Specialist commissioning for managing advanced endometriosis 6

3.2 Specialist service centre aims 8

3.3 CNS responsibilities 8

4 CNS in endometriosis role: skills and knowledge 10

4.1 Clinical practice skills 12

4.2 Leadership skills .....

1 Introduction

Endometriosis is a life challenging disorder that affects over 1.5 million women in the UK – 1 in 10 women and those assigned female at birth from puberty to menopause although the impact may be felt for life. It is a complex illness that is often not quickly diagnosed when women first present with symptoms.

Evidence suggests that care can be delayed due to a lack of awareness and

2 Defining endometriosis

Endometriosis is defined as the presence of endometrial-like tissue outside the uterus, which induces a chronic, inflammatory reaction (NHS, 2022). While some women with endometriosis experience painful symptoms and/or infertility, others have no symptoms at all (or may not recognise their symptoms as abnormal). The exact prevalence of endometriosis is unknown however, WHO (March 2023) states Endometriosis affects roughly 10% (190 million) of reproductive age women and girls globally.

Women of any age can have endometriosis however it is rare before the menarche (first menstrual cycle). Teenagers with painful periods, or who faint, collapse or miss school due to their menstrual periods may be demonstrating symptoms that indicate a possible diagnosis of endometriosis.

Some quick facts and figures about endometriosis

- 1 in 10 women and those assigned female at birth of reproductive age (between puberty and menopause) in the UK suffer from endometriosis.
- 10% of women worldwide have endometriosis – that's 190 million worldwide.
- The prevalence of endometriosis in women with infertility is as high as 50%.
- Endometriosis is the second most common gynaecological condition (after fibroids) in the UK.
- Endometriosis health care costs are comparable to other common diseases such as type 2 diabetes, rheumatoid arthritis, and Crohn's disease.
- On average it takes over 8 years from the onset of symptoms to get a diagnosis.
- Endometriosis costs the UK economy £8.2bn a year in treatment, loss of education, work and health care costs.
- The cause of endometriosis is unknown and there is no definite cure, but many different treatment options.

Information provided by Endometriosis UK

3 The role of the clinical nurse specialist (CNS) in endometriosis

The role of the CNS in managing and supporting women with endometriosis has been defined to take account of the need to:

- lead and develop services
- ensure these services are linked with primary care
- support a better understanding of this condition among all nurses coming in contact with women.

Following the commissioning of complex gynaecology services, the care of women with complex endometriosis has been highlighted. Whilst minor and moderate cases of endometriosis can be managed in all gynaecology departments (with a specialist nurse who is interested in endometriosis), this service specification concerns only severe endometriosis, which has an annual incidence of around 5,000 new cases in the UK per year (NHS England, 2013). However, the RCN supports the need for all women with endometriosis to have access to specialist services to ensure the highest quality care which has been recognised in the NICE guidelines (NICE, 2024). However, this may be difficult to achieve in practice as centres tend to work to hours needed for each complex patient rather than focusing on the needs of individual women, regardless of their situation.

This document builds on this requirement and describes what a CNS might do, and encompasses the care of all women with endometriosis.

To become an endometriosis CNS, nurses will have had:

- extensive experience working within a gynaecology or women's health setting
- will have been educated to masters level and display masters level thinking and decision making
- have insight into the condition and all of the areas of management, including the wider social political dimensions of this condition.

3.1 Specialist commissioning for managing advanced endometriosis

For the purposes of specialist commissioning services, severe endometriosis is defined as either deeply infiltrating endometriosis or recto-vaginal endometriosis.

Deeply infiltrating endometriosis exists where the disease invades at least 5mm below the tissue surface and can occur in a variety of sites; this includes the bladder, pelvic sidewalls, ovaries, pelvic brim, bowel surface and diaphragm.

Recto-vaginal endometriosis is endometriosis which involves the recto-vaginal septum area (recto-vaginal septum; vagina; uterosacral ligaments; rectum). There are many classification systems for endometriosis but none are universally accepted.

Removing the endometriosis involves complex surgery and national BSGE criteria (see bsge.org.uk) exist which set out the standards of service and workload required to undertake surgical excision of severe endometriosis.

This has driven the establishment of centres where such work can be undertaken by specialist multidisciplinary teams. Within the commissioning documents reference is made to an endometriosis CNS whose role is to liaise directly with women using the specialist service and provide women with support in the management of the condition.

The RCN supports NICE guidance (2024) that all women with suspected or confirmed

3.2 Specialist service centre aims

Ky-c32 4(o:- /Span ⁄ET⁄Lang (en-GB)/MCID 259 ⁄BDC680 0 0 1 BT c 0005 11 00005 1w 11 k 0 11 7 0.661056

- careful scrutiny of the referrals by the endometriosis specialist nurse will optimise this arrangement
- women will have contact details of the endometriosis specialist nurse and make contact if problems develop:
  - at six months the endometriosis nurse will review the woman and obtain a completed quality of life questionnaire
  - the same questionnaire will be completed at 12 and 24 months post-surgery and mechanisms for non-face to face consultations (telephone or website submission) need to be in place.

4 Clinical nurse specialist in endometriosis role: skills and knowledge

The role of the CNS endometriosis is complex and will demand a range of practice skills, alongside management and leadership insightfulness.

This senior role will require the nurse to be able to:

•